

Solving the Darwin problem in the first post-Newtonian approximation of general relativity: compressible model

Masaru Shibata* and Keisuke Taniguchi†

**Department of Earth and Space Science, Graduate School of Science, Osaka University,
Toyonaka, Osaka 560, Japan*

†Department of Physics, Kyoto University, Kyoto 606-01, Japan

Abstract

Using the ellipsoidal model for the density configuration, we calculate the equilibrium sequence of the corotating binary stars of the polytropic equation of state in the first post-Newtonian approximation of general relativity. After we calibrate this model by comparing with previous numerical results, we perform the stability analysis by calculating the energy and the angular momentum of the system as a function of the orbital separation. We find that the orbital angular velocity at the energy and/or momentum minimum increases with the increase of the compactness of each star, and this fact holds irrespective of the polytropic index. These features agree with those in previous numerical works. We also show that due to the influence of the tidal field from the companion star, the central density of each star slightly decreases.

PACS number(s): 04.30.Db, 04.25.Nx, 04.40.Dg

I. INTRODUCTION

In a previous paper [1], we analytically obtain the equilibrium configuration of corotating binary stars(i.e., the solution of the Darwin problem) of incompressible fluid in the first post-Newtonian(PN) approximation. We calculate the energy and angular momentum as a function of the orbital separation, and investigate the location of their minimums(we call the orbit at the minimum the innermost stable corotating circular orbit(ISCCO)*). It is found that the angular velocity at the ISCCO increases with the increase of the compactness of the star; i.e., the general relativistic(GR) effect makes the angular velocity at the ISCCO larger.

In this paper, we try to extend the previous work to the case of the compressible fluid star using the ellipsoidal model for the density profile(the so called ellipsoidal approximation) proposed by Lai, Rasio and Shapiro [2]. The ellipsoidal approximation may be used for stars of the sufficiently stiff equation of state(EOS) of a large adiabatic index. Thus, this method may make it possible to investigate the ISCCO for realistic binary neutron stars which are made of the stiff EOS of the large adiabatic index $\Gamma \sim 2 - 3$ [3]. In previous papers, one of us(M.S.) [4] numerically calculates the equilibrium state of binary neutron stars of the polytropic EOS with $\Gamma = 2$ and 3 in the PN approximation. We will show that by a simple semi-analytic calculation, the results obtained in previous papers can be qualitatively explained well. Also, we investigate the influence due to the GR tidal effect in binary neutron stars.

In section II, we show the basic equations to derive the angular velocity for the corotating binary stars. In section III, we introduce the ellipsoidal approximation assuming the polytropic EOS. Using the ellipsoidal approximation, in section IV, we calculate the equations

*The ISCCO is the orbit where the secular instability sets in, and different from the innermost stable circular orbit(ISCO) where the dynamical instability sets in. The ISCO will locate inside the ISCCO [2].

for the energy and angular momentum. In section V, we derive the scalar virial relation in the PN approximation, which is used to determine the size of the star. In section VI, we show equilibrium sequences of binary stars for various polytropic indices. Paying attention to the energy and angular momentum as a function of the orbital separation, we investigate the ISCCO. We also show that due to the influence of the PN tidal field from the companion star, the central density of each star slightly decreases. Section VII is devoted to summary.

Throughout this paper, we take the units of $G = 1$, where G is the gravitational constant, and use c as the light velocity. Latin indices $i, j, k, l, \dots$ take 1 to 3, and δ_{ij} denotes the kronecker's delta. M denotes the Newtonian mass for each star of the binary as

$$M = \int \rho d^3x. \quad (1.1)$$

The quadrupole moment and its tracefree part of each star are defined as

$$I_{ii} = \int \rho x_i^2 d^3x, \quad \text{and} \quad \mathcal{I}_{ii} = I_{ii} - \frac{1}{3} \sum_l I_{ll}. \quad (1.2)$$

II. FORMULATION

In this section, we show the basic equations deriving the equations of the angular velocity and the mass for corotating binary stars in the first PN approximation with no restriction to a special model because these quantities can be written in a simple form without specifying a density configuration. To construct the equilibrium sequence and argue the stability, we additionally need the equations for the energy, the angular momentum, and the scalar virial relation in the PN approximation. We derive them after we specify the model for the density configuration because they are not written in a simple form.

We consider corotating binary stars of equal Newtonian masses ($M_1 = M_2 = M$), whose coordinate separation is R , in the first PN approximation. Here, the gauge condition is the standard PN one [5]. We adopt the coordinates where the center mass of a star (star 1) locates at the origin and the other one (star 2) locates at $(x_1, x_2, x_3) = (-R, 0, 0)$. As

for the density configuration, in this section, we only assume that the star has the triplane symmetry with respect to the symmetric plane(i.e., for star 1, $\rho(x, y, z) = \rho(-x, y, z) = \rho(x, -y, z) = \rho(x, y, -z)$). In the following, when the integral symbol appears, it means that we perform the integral inside the star 1.

In deriving the equations for various quantities, we assume $R > a_0 > M/c^2$, where a_0 is the typical size of the star: We calculate the terms up to $O((a_0/R)^3)$ relative to the lowest order for the angular velocity and the angular momentum, and the terms to $O((a_0/R)^4)$ for the energy. Then, the effects by the quadrupole moment of each star to the equilibrium state are taken into account consistently.

We adopt the polytropic EOS as

$$P = K\rho^\Gamma = (\Gamma - 1)\rho\varepsilon, \quad \Gamma = 1 + \frac{1}{n}, \quad (2.1)$$

where P , ρ , ε , n and K are the pressure, the mass density, the specific internal energy, the polytropic index and the polytropic constant, respectively. Then, in the first PN approximation, the integrated form of the Euler equation is written as [6] [4] [7]

$$\begin{aligned} & K'\rho^{\Gamma-1} - \frac{1}{2c^2} \left(K'\rho^{\Gamma-1} \right)^2 \\ &= U - \frac{X_0}{c^2} + \left\{ \frac{\varpi^2}{2} + \frac{1}{c^2} (2\varpi^2 U - X_\Omega + \beta_\varphi) \right\} \Omega^2 + \frac{\varpi^4}{4c^2} \Omega^4 + \text{constant}, \end{aligned} \quad (2.2)$$

where $\varpi^2 = (x_1 + R/2)^2 + x_2^2$ and $K' = K(n+1)$.

The angular velocity is obtained from the first tensor virial(TV) relation as

$$\begin{aligned} 0 = \int \frac{\partial P}{\partial x_1} \left(1 - \frac{\Gamma\varepsilon}{c^2} \right) d^3x &= \int \rho U_{,1} d^3x + \frac{R}{2} M \Omega^2 + \frac{1}{c^2} \left[- \int \rho X_{0,1} d^3x \right. \\ &\quad \left. + \Omega_N^2 \int \rho (2\varpi^2 U_{,1} + 4x_1 U + 2RU - X_{\Omega,1} + \beta_{\varphi,1}) d^3x \right. \\ &\quad \left. + \Omega_N^4 \left(\frac{R}{2} (3I_{11} + I_{22}) + \frac{MR^3}{8} \right) \right], \end{aligned} \quad (2.3)$$

where $\partial/\partial x_k$ denotes the partial derivative with respect to x_k . Ω_N denotes the angular velocity in the Newtonian order solution as [2]

$$\Omega_N^2 = \frac{2M}{R^3} + \frac{18I_{11}}{R^5}. \quad (2.4)$$

To determine the orbital angular velocity up to the PN order, we need to perform the following integrals:

$$I_1 = \int \rho U d^3x, \quad (2.5)$$

$$I_2 = \int \rho U_{,1} d^3x, \quad (2.6)$$

$$I_3 = \int \rho U x_1 d^3x, \quad (2.7)$$

$$I_4 = \int \rho U_{,1} x_1 d^3x, \quad (2.8)$$

$$I_5 = \int \rho U_{,1} x_k^2 d^3x, \quad (2.9)$$

$$I_6 = \int \rho X_{0,1} d^3x, \quad (2.10)$$

$$I_7 = \int \rho X_{\Omega,1} d^3x, \quad (2.11)$$

$$I_8 = \int \rho \beta_{\varphi,1} d^3x. \quad (2.12)$$

In the following subsections, we separately show the equations for the gravitational potentials and their solutions to evaluate $I_1 \sim I_8$.

A. Equation and solution for U and integrals including U : $I_1 \sim I_5$

Equation for the Newtonian potential U is

$$\Delta U = -4\pi\rho. \quad (2.13)$$

Here, there are two kinds of contribution for U at star 1; one is the contribution from star 1 itself and the other is that from star 2. Hence, we write U as

$$U = U^{1 \rightarrow 1} + U^{2 \rightarrow 1}, \quad (2.14)$$

where $U^{i \rightarrow j}$ denotes the potential generated by star i at star j . Here, $U^{2 \rightarrow 1}$ is generally expressed as

$$\begin{aligned} U^{2 \rightarrow 1} = & \frac{M}{R} \left(1 - \frac{x_1}{R} + \frac{2x_1^2 - x_2^2 - x_3^2}{2R^2} + \frac{-2x_1^3 + 3x_1(x_2^2 + x_3^2)}{2R^3} + O(R^{-4}) \right) \\ & + \frac{3I_{11}}{2R^3} \left(1 - \frac{3x_1}{R} + O(R^{-2}) \right). \end{aligned} \quad (2.15)$$

On the other hand, we cannot specify $U^{1\rightarrow 1}$ without the density configuration. Thus, we define the following integrals [8];

$$W = \frac{1}{2} \int \rho U^{1\rightarrow 1} d^3x, \quad (2.16)$$

$$\mathcal{M}_{ii} = \int \rho U_{,i}^{1\rightarrow 1} x_i d^3x, \quad (2.17)$$

where we have the relation $\sum_i \mathcal{M}_{ii} = -W$ [8]. Using the solution for U , we get

$$I_1 = 2W + \frac{M^2}{R} + \frac{3M}{R^3} \mathcal{I}_{11} + O(R^{-5}), \quad (2.18)$$

$$I_2 = -\frac{M^2}{R^2} - \frac{9M \mathcal{I}_{11}}{R^4} + O(R^{-6}), \quad (2.19)$$

$$I_3 = -\frac{M}{R^2} I_{11} + O(R^{-4}), \quad (2.20)$$

$$I_4 = \mathcal{M}_{11} + \frac{2M}{R^3} I_{11} + O(R^{-5}), \quad (2.21)$$

$$I_5 = -\frac{M}{R^2} I_{kk} + O(R^{-4}). \quad (2.22)$$

Note that Ω_N is derived using I_2 .

B. Equation and solution for X_0 and integrals including X_0 : I_6

As in the case of U , X_0 is divided into two parts as $X_0 = X_0^{1\rightarrow 1} + X_0^{2\rightarrow 1}$. We also write I_6 as $I_6^{1\rightarrow 1} + I_6^{2\rightarrow 1}$, where $I_6^{1\rightarrow 1}$ and $I_6^{2\rightarrow 1}$ correspond to the contributions from $X_0^{1\rightarrow 1}$ and $X_0^{2\rightarrow 1}$, respectively. Equations for $X_0^{1\rightarrow 1}$ and $X_0^{2\rightarrow 1}$ are

$$\Delta X_0^{1\rightarrow 1} = 4\pi\rho \left[2U^{1\rightarrow 1} + 2U^{2\rightarrow 1} + \varepsilon + \frac{3P}{\rho} \right] \equiv 4\pi\rho_X^{1\rightarrow 1}, \quad (2.23)$$

$$\Delta X_0^{2\rightarrow 1} = 4\pi\rho \left[2U^{2\rightarrow 2} + 2U^{1\rightarrow 2} + \varepsilon + \frac{3P}{\rho} \right] \equiv 4\pi\rho_X^{2\rightarrow 1}. \quad (2.24)$$

To evaluate $I_6^{1\rightarrow 1}$, we need only the odd function of x_1 in $\rho_X^{1\rightarrow 1}$, and the terms required are

$$\rho_X^{1\rightarrow 1} = -2\rho \frac{x_1}{R^2} \left(M + M \frac{2x_1^2 - 3x_2^2 - 3x_3^2}{2R^2} + \frac{9\mathcal{I}_{11}}{2R^2} + O(R^{-4}) \right). \quad (2.25)$$

Then, the terms of $X_0^{1\rightarrow 1}$ required for $I_6^{1\rightarrow 1}$ are

$$X_0^{1\rightarrow 1} = \left(\frac{2M}{R^2} + \frac{9\mathcal{I}_{11}}{R^4} \right) D_1 + \frac{M}{R^4} (2D_{111} - 3D_{122} - 3D_{133}) + O(R^{-6}), \quad (2.26)$$

where D_i and D_{1kk} are the solutions of

$$\Delta D_i = -4\pi\rho x_i, \quad (2.27)$$

$$\Delta D_{1kk} = -4\pi\rho x_1 x_k^2. \quad (2.28)$$

Using the relations,

$$\int \rho D_{i,i} d^3x = -\mathcal{M}_{ii}, \quad (2.29)$$

$$\int \rho D_{1kk,1} d^3x = -\int \rho U_{,1}^{1\rightarrow 1} x_1 x_k^2 d^3x \equiv -\mathcal{M}_{11kk}, \quad (2.30)$$

$I_6^{1\rightarrow 1}$ is evaluated to be

$$I_6^{1\rightarrow 1} = -\left(\frac{2M}{R^2} + \frac{9\mathcal{I}_{11}}{R^4}\right)\mathcal{M}_{11} - \frac{M}{R^4}(2\mathcal{M}_{1111} - 3\mathcal{M}_{1122} - 3\mathcal{M}_{1133}) + O(R^{-6}). \quad (2.31)$$

On the other hand, the solution of $X_0^{2\rightarrow 1}$ is written as

$$\begin{aligned} X_0^{2\rightarrow 1} = & \frac{M_X}{R} \left(1 - \frac{x_1}{R} + \frac{2x_1^2 - x_2^2 - x_3^2}{2R^2} + \frac{-2x_1^3 + 3x_1x_2^2 + 3x_1x_3^2}{2R^3} + O(R^{-4})\right) \\ & + \frac{d_X}{R^2} \left(1 - \frac{2x_1}{R} + O(R^{-2})\right) + \frac{3\mathcal{I}_{X11}}{2R^3} \left(1 - \frac{3x_1}{R} + O(R^{-2})\right), \end{aligned} \quad (2.32)$$

where

$$M_X = -\int \rho_X^{2\rightarrow 1} d^3x = -4W - (3\Gamma - 2)U_i - 2\left(\frac{M^2}{R} + \frac{3M\mathcal{I}_{11}}{R^3}\right) + O(R^{-5}), \quad (2.33)$$

$$d_X = -\int \rho_X^{2\rightarrow 1} x_1 d^3x = -\frac{2MI_{11}}{R^2} + O(R^{-4}), \quad (2.34)$$

$$\begin{aligned} \mathcal{I}_{X11} = & -\frac{1}{3} \int \rho_X^{2\rightarrow 1} (2x_1^2 - x_2^2 - x_3^2) d^3x \\ = & -2\mathcal{I}_{U11} - (3\Gamma - 2)\mathcal{I}_{\varepsilon 11} - \frac{2M}{R}\mathcal{I}_{11} + O(R^{-3}), \end{aligned} \quad (2.35)$$

and

$$U_i = \int \rho \varepsilon d^3x, \quad (2.36)$$

$$\mathcal{I}_{U11} = \frac{2I_{U11} - I_{U22} - I_{U33}}{3}, \quad I_{Ukk} \equiv \int \rho U^{1\rightarrow 1} x_k^2 d^3x, \quad (2.37)$$

$$\mathcal{I}_{\varepsilon 11} = \frac{2I_{\varepsilon 11} - I_{\varepsilon 22} - I_{\varepsilon 33}}{3}, \quad I_{\varepsilon kk} \equiv \int \rho \varepsilon x_k^2 d^3x. \quad (2.38)$$

Thus, $I_6^{2\rightarrow 1}$ is evaluated to be

$$\begin{aligned}
I_6^{2 \rightarrow 1} &= -M \left(\frac{M_X}{R^2} + \frac{2d_X}{R^3} + \frac{9M_X \mathcal{I}_{11}}{2MR^4} + \frac{9\mathcal{I}_{X11}}{2R^4} \right) \\
&= M \left[\frac{4W + (3\Gamma - 2)U_i}{R^2} + \frac{2M^2}{R^3} + \frac{24M \mathcal{I}_{11}}{R^5} + \frac{4MI_{11}}{R^5} \right. \\
&\quad \left. + \frac{9}{2R^4} \left\{ (4W + (3\Gamma - 2)U_i) \frac{\mathcal{I}_{11}}{M} + 2\mathcal{I}_{U11} + (3\Gamma - 2)\mathcal{I}_{\varepsilon 11} \right\} + O(R^{-6}) \right]. \quad (2.39)
\end{aligned}$$

C. Equation and solution for X_Ω and integral including X_Ω : I_7

Equation for X_Ω is

$$\Delta X_\Omega = 8\pi\rho \left(x_1^2 + x_2^2 + Rx_1 + \frac{R^2}{4} \right). \quad (2.40)$$

The solution is also written as $X_\Omega^{1 \rightarrow 1} + X_\Omega^{2 \rightarrow 1}$, where

$$X_\Omega^{1 \rightarrow 1} = -2 \left(D_{11} + D_{22} + RD_1 + \frac{R^2}{4} U^{1 \rightarrow 1} \right), \quad (2.41)$$

$$X_\Omega^{2 \rightarrow 1} = -\frac{R^2}{2} U^{2 \rightarrow 1} + 2 \frac{I_{11}}{R} \left(1 - \frac{2x_1}{R} + O(R^{-2}) \right) - \frac{2(I_{11} + I_{22})}{R} \left(1 - \frac{x_1}{R} + O(R^{-2}) \right), \quad (2.42)$$

and D_{kk} is the solution for $\Delta D_{kk} = -4\pi\rho x_k^2$. Using these solutions, we soon get I_7 as

$$I_7 = 2R\mathcal{M}_{11} + \frac{M}{2} \left(M + \frac{9\mathcal{I}_{11}}{R^2} - \frac{4}{R^2} (I_{11} - I_{22}) + O(R^{-3}) \right). \quad (2.43)$$

D. Equation and solution for β_φ and integral including β_φ : I_8

Definition of β_φ is

$$\begin{aligned}
\beta_\varphi &= -\frac{7}{2} \left(x_1 P_1 + x_2 P_2 + \frac{R}{2} P_1 \right) \\
&\quad - \frac{1}{2} \left[\left(x_1 + \frac{R}{2} \right)^2 P_{2,2} + x_2^2 P_{1,1} - \left(x_1 + \frac{R}{2} \right) x_2 (P_{1,2} + P_{2,1}) \right], \quad (2.44)
\end{aligned}$$

where

$$\Delta P_1 = -4\pi\rho \left(x_1 + \frac{R}{2} \right), \quad (2.45)$$

$$\Delta P_2 = -4\pi\rho x_2. \quad (2.46)$$

P_i is also written as $P_i^{1\rightarrow 1} + P_i^{2\rightarrow 1}$, where

$$P_1^{1\rightarrow 1} = D_1 + \frac{R}{2}U^{1\rightarrow 1}, \quad (2.47)$$

$$P_2^{1\rightarrow 1} = D_2, \quad (2.48)$$

and

$$P_1^{2\rightarrow 1} = \frac{I_{11}}{R^2} \left(1 - \frac{2x_1}{R} + \frac{6x_1^2 - 3x_2^2 - 3x_3^2}{2R^2} + O(R^{-3}) \right) - \frac{R}{2}U^{2\rightarrow 1}, \quad (2.49)$$

$$P_2^{2\rightarrow 1} = \frac{I_{22}}{R^2} \left(\frac{x_2}{R} - \frac{3x_1x_2}{R^2} + O(R^{-3}) \right). \quad (2.50)$$

$\beta_\varphi^{2\rightarrow 1}$ can be written to the explicit form as

$$\begin{aligned} \beta_\varphi^{2\rightarrow 1} = & \frac{MR}{8} \left(7 + \frac{7x_1}{R} - \frac{14x_1^2 + 9x_2^2 + 7x_3^2}{2R^2} + \frac{x_1(14x_1^2 + 13x_2^2 + 7x_3^2)}{2R^3} + O(R^{-4}) \right) \\ & + \frac{21I_{11}}{16R} \left(1 - \frac{x_1}{R} + O(R^{-2}) \right) - \frac{7I_{11}}{4R} \left(1 + O(R^{-2}) \right) - \frac{I_{22}}{8R} \left(1 + \frac{x_1}{R} + O(R^{-2}) \right). \end{aligned} \quad (2.51)$$

Then, after straightforward calculations, I_8 is evaluated as

$$I_8 = -\frac{R}{2}(7W - \mathcal{M}_{22}) + \frac{7M^2}{8} + \frac{M}{8R^2}(14I_{11} + 9I_{22} + 7I_{33}) + O(R^{-3}). \quad (2.52)$$

E. Ω^2

Gathering the results obtained in previous subsections, Ω^2 is calculated as

$$\begin{aligned} \Omega^2 = & \frac{2M}{R^3} \left[1 + \frac{1}{Mc^2} \left\{ 4W + (3\Gamma - 2)U_i + \mathcal{M}_{33} - \mathcal{M}_{11} - \frac{9M^2}{4R} \right. \right. \\ & \left. \left. - \frac{M}{2R^3} (28I_{11} - 16I_{22} - 5I_{33}) + O(R^{-4}) \right\} \right] \\ & + \frac{18I_{11}}{R^5} \left[1 + \frac{1}{c^2M} \left(W - \mathcal{M}_{11} - \mathcal{M}_{22} + \frac{3\Gamma - 2}{2}U_i + O(R^{-2}) \right) \right] \\ & + \frac{1}{c^2R^5} \left(18I_{U11} + 9(3\Gamma - 2)I_{\varepsilon 11} - 4\mathcal{M}_{1111} + 6\mathcal{M}_{1122} + 6\mathcal{M}_{1133} \right). \end{aligned} \quad (2.53)$$

To transform this equation into a more simple form, we use the scalar, the second and the fourth tensor virial equations in the Newtonian order as follows;

$$3(\Gamma - 1)U_i = W - \frac{3M}{R^3}I_{11} - \Omega_N^2(I_{11} + I_{22}) + O(R^{-5}), \quad (2.54)$$

$$\mathcal{M}_{11} = -(\Gamma - 1)U_i - \Omega_N^2 I_{11} - \frac{2MI_{11}}{R^3} + O(R^{-5}), \quad (2.55)$$

$$\mathcal{M}_{22} = -(\Gamma - 1)U_i - \Omega_N^2 I_{22} + \frac{MI_{22}}{R^3} + O(R^{-5}), \quad (2.56)$$

$$\mathcal{M}_{33} = -(\Gamma - 1)U_i + \frac{MI_{33}}{R^3} + O(R^{-5}), \quad (2.57)$$

$$2\mathcal{M}_{1111} - 3\mathcal{M}_{1122} - 3\mathcal{M}_{1133} = -9(\Gamma - 1)I_{\varepsilon 11} + O(R^{-3}). \quad (2.58)$$

Then,

$$\begin{aligned} \Omega^2 = & \frac{2M}{R^3} \left[1 + \frac{1}{Mc^2} \left\{ 5W + U_i - \frac{9M^2}{4R} - \frac{M}{2R^3} (28I_{11} - 14I_{22} - 9I_{33}) + O(R^{-4}) \right\} \right] \\ & + \frac{18I_{11}}{R^5} \left[1 + \frac{1}{c^2} \left(\frac{13+n}{6} \frac{W}{M} + O(R^{-2}) \right) \right] + \frac{1}{c^2 R^5} (18I_{U11} + 9(5\Gamma - 4)I_{\varepsilon 11}). \end{aligned} \quad (2.59)$$

We note that in the incompressible case ($\rho = \text{constant}$ and $\Gamma \rightarrow \infty$), the configuration of each star is the ellipsoidal figure of its axial length a_1 , a_2 , and a_3 , and we have relations

$$W = \frac{2M\pi\rho A_0}{5}, \quad (2.60)$$

$$U_i = 0, \quad (2.61)$$

$$I_{U11} = \frac{16\pi\rho A_0}{21} I_{11}, \quad (2.62)$$

$$(\Gamma - 1)I_{\varepsilon 11} = \frac{2\pi\rho A_0}{21} I_{11}, \quad (2.63)$$

where [8]

$$\begin{aligned} A_0 &= a_1 a_2 a_3 \int_0^\infty \frac{du}{\sqrt{(a_1^2 + u)(a_2^2 + u)(a_3^2 + u)}} \\ &= a_1^2 \alpha_2 \alpha_3 \int_0^\infty \frac{dt}{\sqrt{(1 + t)(\alpha_2^2 + t)(\alpha_3^2 + t)}} \equiv a_1^2 \hat{A}_0(\alpha_2, \alpha_3), \end{aligned} \quad (2.64)$$

and $\alpha_2 = a_2/a_1$ and $\alpha_3 = a_3/a_1$. Then, Ω^2 reduces to that derived in a previous paper [1].

F. Mass and center of mass

The conserved mass is defined as [5] [4]

$$\begin{aligned}
M_* &= \int d^3x \rho_* = \int d^3x \rho \left[1 + \frac{1}{c^2} \left(\frac{v^2}{2} + 3U \right) \right] \\
&= M \left[1 + \frac{1}{c^2} \left(\frac{6W}{M} + \frac{13M}{4R} + \frac{1}{4R^3} (45I_{11} + 4I_s) + O(R^{-5}) \right) \right],
\end{aligned} \tag{2.65}$$

where $I_s = I_{11} + I_{22}$. The PPN mass [9] is slightly different from M_* as

$$\begin{aligned}
M_{\text{PPN}} &= \int d^3x \rho \left[1 + \frac{1}{c^2} \left\{ \frac{v^2}{2} + \varepsilon + 3U - \frac{U^{1 \rightarrow 1}}{2} + \frac{\Omega_N^2}{2} (x_1^2 + x_2^2) \right\} \right] \\
&= M \left[1 + \frac{1}{c^2} \left(\frac{5W + U_i}{M} + \frac{13M}{4R} + \frac{1}{4R^3} (45I_{11} + 8I_s) + O(R^{-5}) \right) \right].
\end{aligned} \tag{2.66}$$

The advantage of the PPN mass is that if we use it, Ω^2 can be denoted only by M_{PPN} for the limit $I_{kk}/R^3 \rightarrow 0$ [9]. However, when I_{kk} is large enough, it is not the conserved quantity any more, and not so useful.

We define the center of mass by the conserved mass density as

$$x_*^i = \frac{1}{M_*} \int d^3x \rho_* x^i, \tag{2.67}$$

and the x_1 coordinate of the center of mass for star 1 becomes

$$\frac{1}{c^2} \left(-\frac{2I_{11}}{R^2} + O(R^{-4}) \right). \tag{2.68}$$

Note that in the PPN formalism [9], the center of mass of each star is defined as

$$x_{\text{PPN}}^i = \frac{1}{M_{\text{PPN}}} \int d^3x \rho x^i \left[1 + \frac{1}{c^2} \left\{ \frac{v^2}{2} + \varepsilon + 3U - \frac{U^{1 \rightarrow 1}}{2} + \frac{\Omega_N^2}{2} (x_1^2 + x_2^2) \right\} \right], \tag{2.69}$$

and the results are the same. From Eq.(2.68) we define the orbital separation as

$$R_* = R_{\text{PPN}} = R \left\{ 1 + \frac{1}{c^2} \left(-\frac{4I_{11}}{R^3} + O(R^{-5}) \right) \right\}. \tag{2.70}$$

III. THE ELLIPSOIDAL APPROXIMATION

In this section, we introduce the ellipsoidal approximation for the density configuration proposed by Lai, Rasio and Shapiro(LRS) [2]. In this approximation, the configuration of each star is assumed to be the ellipsoidal figure of its axial length a_1 , a_2 and a_3 . Then, the density profile is assumed to be

$$\rho = \rho_p(\xi), \quad \text{for } \xi = \sqrt{\frac{x_1^2}{\hat{a}_1^2} + \frac{x_2^2}{\hat{a}_2^2} + \frac{x_3^2}{\hat{a}_3^2}} \leq \xi_1 \quad \text{and} \quad \hat{a}_k = \frac{a_k}{\xi_1}, \quad (3.1)$$

where $\rho_p(\xi)$ has the density profile of the spherical polytrope, and $\rho = 0$ for $\xi > \xi_1$. Because of the polytropic relation, P and ε are also assumed to depend only on ξ .

Here, there is no solid basis to justify this model: i.e., in this approximation, we do not rigidly treat (a) the change of the density profile due to the PN gravity which occurs even for the spherical star, and (b) the deformation of the star due to the tidal gravity by the companion star. These facts mean that several quantities of the system calculated below such as the energy, the angular momentum, the central density of the star, and so on, deviate systematically from their exact values (see section VI). However, (1) in the Newtonian case, this treatment is almost exact for the incompressible fluid [8] [2], and (2) in a previous paper [1], we find that the tidal deformation by the PN gravity is a small effect for the incompressible fluid star. Thus, we may expect that the ellipsoidal approximation is good for a sufficiently stiff EOS (small n).

Following the standard procedure, we set

$$\rho_p = \rho_c \theta(\xi)^n, \quad (3.2)$$

where ρ_c is the central density, and θ obeys the Lane-Emden equation [3]. Then, M , I_{kk} , U_i , and $I_{\varepsilon kk}$ are immediately calculated as

$$M = 4\pi\rho_c a_1 a_2 a_3 \xi_1^{-1} |\theta'|, \quad (3.3)$$

$$I_{kk} = \frac{M a_k^2}{5} \kappa_n, \quad \kappa_n \equiv \frac{5}{3\xi_1^4 |\theta'|} \int_0^{\xi_1} \theta^n \xi^4 d\xi, \quad (3.4)$$

$$U_i = k_1 K \rho_c^{1/n} M, \quad k_1 \equiv \frac{n(n+1)}{5-n} \xi_1 |\theta'|, \quad (3.5)$$

$$I_{\varepsilon kk} = 5nK \rho_c^{1/n} I_{kk} \frac{\sigma_n}{\kappa_n}, \quad \sigma_n \equiv \frac{1}{3\xi_1^4 |\theta'|} \int_0^{\xi_1} \theta^{n+1} \xi^4 d\xi, \quad (3.6)$$

where $\theta' = \partial\theta/\partial\xi$ at ξ_1 .

As for W and $\mathcal{M}_{kk}$, we follow LRS [2] and set as

$$W = \frac{2\pi\bar{\rho}A_0}{5-n} M, \quad (3.7)$$

$$\mathcal{M}_{kk} = -\frac{2\pi\bar{\rho}A_k a_k^2}{5-n} M, \quad (3.8)$$

where $\bar{\rho} = 3M/(4\pi a_1 a_2 a_3)$, in order to guarantee that in the case of the incompressible ellipsoid or the spherical star, these are exact definitions. Due to the same principle, I_{Ukk} is written as

$$I_{Ukk} = \frac{5-n}{3} \frac{W}{M} I_{kk} \left(\frac{5\sigma_n}{\xi_1 |\theta'| \kappa_n} + 1 \right), \quad (3.9)$$

where we make use of the fact that in the case of the spherical star and/or the incompressible star, $U^{1 \rightarrow 1}$ is written as

$$U^{1 \rightarrow 1} = \frac{5-n}{3} \frac{W}{M} \left(1 + \frac{\theta}{\xi_1 |\theta'|} \right). \quad (3.10)$$

Thus, all quantities to calculate Ω^2 and M_* reduce to the available form, and they are rewritten as

$$\begin{aligned} \Omega^2 &= \frac{M}{a_1^3} \left(\frac{2}{\hat{R}^3} + \frac{6\kappa_n q_{11}}{5\hat{R}^5} \right) + \frac{M^2}{c^2 a_1^4} \left[\frac{2}{\hat{R}^3} \left\{ 5\hat{W} + \hat{U}_i - \frac{9}{4\hat{R}} - \frac{\kappa_n}{10\hat{R}^3} (28 - 14\alpha_2^2 - 9\alpha_3^2) \right\} \right. \\ &\quad \left. + \frac{6\kappa_n q_{11}}{5\hat{R}^5} \hat{W} \left(\frac{23-n}{6} + \gamma_n \frac{15n+35}{6(n+1)} \right) \right] \\ &\equiv \frac{M}{a_1^3} \hat{\Omega}_N^2(\hat{R}) + \frac{M^2}{c^2 a_1^4} \hat{\Omega}_{PN}^2(\hat{R}), \end{aligned} \quad (3.11)$$

$$M_* = M \left[1 + \frac{M}{c^2 a_1} \left\{ 6\hat{W} + \frac{13}{4\hat{R}} + \frac{\kappa_n}{20\hat{R}^3} (15q_{11} + 4q_s) \right\} \right], \quad (3.12)$$

where $q_{11} = 2 - \alpha_2^2 - \alpha_3^2$, $q_s = 1 + \alpha_2^2$, $\hat{R} = R/a_1$, $\gamma_n = (5-n)\sigma_n/(\xi_1 |\theta'| \kappa_n)$, and

$$\hat{W}(\hat{R}) = W \frac{a_1}{M^2} = \frac{3\hat{A}_0}{2(5-n)\alpha_2\alpha_3}, \quad (3.13)$$

$$\hat{U}_i(\hat{R}) = U_i \frac{a_1}{M^2} = \frac{n}{3} \left(\hat{W} - \frac{\kappa_n q_{11}}{5\hat{R}^3} - \frac{\kappa_n q_s}{5} \hat{\Omega}_N^2 \right). \quad (3.14)$$

Here, to derive the expression for $\hat{U}_i$, we use the scalar virial relation in the Newtonian order. In the ellipsoidal approximation, α_2 and α_3 , which are functions of $\hat{R}$, are assumed to be determined from the tensor virial relations in the Newtonian order as [2]

$$\begin{aligned} \frac{2\pi\bar{\rho}A_1a_1^2}{5-n}M &= \frac{1}{n}k_1K\rho_c^{1/n}M + \frac{2MI_{11}}{R^3} + \Omega_N^2 I_{11}, \\ \frac{2\pi\bar{\rho}A_2a_2^2}{5-n}M &= \frac{1}{n}k_1K\rho_c^{1/n}M - \frac{MI_{22}}{R^3} + \Omega_N^2 I_{22}, \\ \frac{2\pi\bar{\rho}A_3a_3^2}{5-n}M &= \frac{1}{n}k_1K\rho_c^{1/n}M - \frac{MI_{33}}{R^3}. \end{aligned} \quad (3.15)$$

IV. THE ENERGY AND THE ANGULAR MOMENTUM

Next, we consider the total energy of the system. The energy for each star is calculated from $E_1 = E_N + E_{\text{PN}}/c^2$, where [5] [4]

$$E_N = \int \rho \left(\varepsilon + \frac{1}{2}v^2 - \frac{1}{2}U \right) d^3x, \quad (4.1)$$

$$E_{\text{PN}} = \int \rho \left(\frac{5}{8}v^4 + \frac{5}{2}v^2U + \Gamma\varepsilon v^2 + 2\varepsilon U - \frac{5}{2}U^2 + \frac{1}{2}\beta_\varphi\Omega^2 \right) d^3x, \quad (4.2)$$

and $v^2 = \varpi^2\Omega^2$. Even for the compressible PN Darwin problem, the expression of E_N is soon derived by using the quantities defined above as

$$\begin{aligned} E_N &= U_i - W + \frac{\Omega^2}{2} \left(\frac{MR^2}{4} + I_s \right) - \frac{M}{2} \left(\frac{M}{R} + \frac{3\mathcal{I}_{11}}{R^3} \right) \\ &= \left(\frac{M^2}{a_1} \right) \left[\hat{U}_i - \hat{W} + \frac{\hat{\Omega}^2}{2} \left(\frac{\hat{R}^2}{4} + \frac{\kappa_n q_s}{5} \right) - \frac{1}{2} \left(\frac{1}{\hat{R}} + \frac{\kappa_n q_{11}}{5\hat{R}^3} \right) \right] \\ &\equiv \left(\frac{M^2}{a_1} \right) \left[\hat{E}_N(\hat{R}) + \frac{M}{2a_1c^2} \hat{\Omega}_{\text{PN}}^2 \left(\frac{\hat{R}^2}{4} + \frac{\kappa_n q_s}{5} \right) \right], \end{aligned} \quad (4.3)$$

where $\hat{\Omega}^2 = \Omega^2 a_1^3/M = \hat{\Omega}_N^2 + (M/c^2 a_1) \hat{\Omega}_{\text{PN}}^2$. On the other hand, to express E_{PN} , we are required to evaluate new integrals as

$$J_1 = \int \rho \varepsilon U^{1 \rightarrow 1} d^3x, \quad (4.4)$$

$$J_2 = \int \rho (U^{1 \rightarrow 1})^2 d^3x, \quad (4.5)$$

$$J_3 = \int \rho \varpi^2 U^{1 \rightarrow 1} d^3x, \quad (4.6)$$

$$J_4 = \int \rho U^{1 \rightarrow 1} U^{2 \rightarrow 1} d^3x, \quad (4.7)$$

$$J_5 = \int \rho \beta_\varphi^{1 \rightarrow 1} d^3x. \quad (4.8)$$

To evaluate $J_1 \sim J_4$, we use the same principle in calculating $\mathcal{I}_{U11}$. Then, they become

$$J_1 = \frac{n}{3} K \rho_c^{1/n} W \xi_1 |\theta'| \left(\frac{(5-n)\lambda_n}{\xi_1^2 |\theta'|^2} + n + 1 \right), \quad (4.9)$$

$$J_2 = \frac{5-n}{9} \frac{W^2}{M} \left(\frac{(5-n)\lambda_n}{\xi_1^2 |\theta'|^2} + n + 7 \right), \quad (4.10)$$

$$J_3 = \frac{R^2 W}{2} + \frac{5-n}{3} \frac{W}{M} I_s \left(5 \frac{\sigma_n}{\xi_1 |\theta'| \kappa_n} + 1 \right), \quad (4.11)$$

$$J_4 = 2W \left(\frac{M}{R} + \frac{3\mathcal{I}_{11}}{2R^3} \right) + \frac{\mathcal{I}_{11}}{2R^3} (5-n) W \left(5 \frac{\sigma_n}{\xi_1 |\theta'| \kappa_n} + 1 \right) + O(R^{-5}), \quad (4.12)$$

where

$$\lambda_n = \frac{1}{\xi_1^2 |\theta'|} \int_0^{\xi_1} \theta^{n+2} \xi^2 d\xi. \quad (4.13)$$

On the other hand, J_5 is written as

$$\begin{aligned} J_5 = & \frac{R^2}{4} (\mathcal{M}_{22} - 7W) - \frac{7}{2} \int \rho (D_1 x_1 + D_2 x_2) d^3 x \\ & - \frac{1}{2} \int \rho (x_2^2 D_{1,1} + x_1^2 D_{2,2} - x_1 x_2 D_{1,2} - x_1 x_2 D_{2,1}) d^3 x. \end{aligned} \quad (4.14)$$

Unfortunately, the integrand in the second and third terms cannot be expressed by θ . Hence, we need to introduce a new function: To evaluate D_1 and D_2 , we assume that the density profile ρ , which appears in the source terms for their Poisson equations, is equal to that of the spherical polytrope $\rho_p(\xi)$. Then, we can write D_k as

$$D_k = 4\pi \rho_c \hat{a}^2 \phi(\xi) x_k, \quad (4.15)$$

where $\hat{a}^3 = \hat{a}_1 \hat{a}_2 \hat{a}_3$ and ϕ satisfies

$$\frac{d^2 \phi}{d\xi^2} + \frac{4}{\xi} \frac{d\phi}{d\xi} = -\theta^n. \quad (4.16)$$

Note that the following relation holds;

$$\sum_k D_{k,k} = U^{1 \rightarrow 1} + \sum_k x_k U_{,k}^{1 \rightarrow 1}. \quad (4.17)$$

Also, for the spherical star,

$$4\pi \rho_c \hat{a}^2 = \frac{5-n}{3\xi_1 |\theta'|} \frac{W}{M}. \quad (4.18)$$

From these two equations, we have the relation

$$3\phi + \xi \frac{d\phi}{d\xi} = \xi_1 |\theta'| + \theta + \xi \frac{d\theta}{d\xi}. \quad (4.19)$$

Thus, ϕ is solved under the boundary conditions $\phi = (\xi_1 |\theta'| + 1)/3$ and $d\phi/d\xi = 0$ at $\xi = 0$.

Using this definition of D_k , J_5 can be written as

$$J_5 = \frac{R^2}{4} (\mathcal{M}_{22} - 7W) - \frac{4M^2}{3(a_1 a_2 a_3)^{1/3}} (a_1^2 + a_2^2) \eta_n, \quad (4.20)$$

where

$$\eta_n = \frac{1}{\xi_1^5 |\theta'|^2} \int_0^{\xi_1} \theta^n \phi \xi^4 d\xi. \quad (4.21)$$

Exchanging $M^2/(a_1 a_2 a_3)^{1/3}$ into $(5-n)W/3$, we finally get

$$J_5 = \frac{R^2}{4}(\mathcal{M}_{22} - 7W) - \frac{4W}{9}(5-n)(a_1^2 + a_2^2)\eta_n. \quad (4.22)$$

Gathering the above results, we obtain the expression for E_{PN} as

$$\begin{aligned} E_{\text{PN}} = & -\frac{5(5-n)}{18} \frac{W^2}{M} (\tilde{\lambda}_n + n + 7) + \frac{2(5-n)}{3(n+1)} \frac{U_i W}{M} (\tilde{\lambda}_n + n + 1) \\ & + \frac{M}{R} \left\{ -\frac{37}{4} W + \left(\frac{5n}{2} + \frac{1}{2} \right) \frac{U_i}{n} + \frac{\mathcal{M}_{22}}{4} \right\} - \frac{7M^3}{32R^2} \\ & + \frac{1}{R^3} \left[W \left\{ -\mathcal{I}_{11} \left(\frac{83-10n}{4} + \frac{25}{2} \gamma_n \right) + I_s \left(\frac{25}{3} \gamma_n + \frac{5(5-n)}{3} - \frac{20(5-n)}{9} \frac{\eta_n}{\kappa_n} \right) \right\} \right. \\ & \quad \left. + \frac{9}{4} \mathcal{I}_{11} \mathcal{M}_{22} + \frac{U_i}{n} \left\{ \left(\frac{15n}{2} + \frac{9}{2} \right) \mathcal{I}_{11} + \frac{5\gamma_n}{n+1} (2(n+1)I_s + 3n\mathcal{I}_{11}) \right\} \right] \\ & + \frac{M^2}{16R^4} (106I_{11} + 25I_{22} - 71I_{33}), \end{aligned} \quad (4.23)$$

where $\tilde{\lambda}_n = (5-n)\lambda_n/(\xi_1^2 |\theta'|^2)$. Using the scalar and tensor virial relations in the Newtonian order, E_{PN} is rewritten to a simple form

$$\begin{aligned} E_{\text{PN}} = & -\frac{5(5-n)}{18} \frac{W^2}{M} (\tilde{\lambda}_n + n + 7) + \frac{2(5-n)}{3(n+1)} \frac{U_i W}{M} (\tilde{\lambda}_n + n + 1) \\ & + \frac{M}{R} \left(-\frac{55}{6} W + \frac{5}{2} U_i \right) - \frac{7M^3}{32R^2} \\ & + \frac{W}{R^3} \left[\mathcal{I}_{11} \left(5n - 20 + \frac{5n}{n+1} \gamma_n - \frac{25}{2} \gamma_n \right) + I_s \left(\frac{35}{3} \gamma_n + \frac{5(5-n)}{3} - \frac{20(5-n)}{9} \frac{\eta_n}{\kappa_n} \right) \right] \\ & + \frac{M^2}{48R^4} (302I_{11} + 59I_{22} - 209I_{33}) + O(R^{-5}) \\ \equiv & \left(\frac{M^3}{a_1^2} \right) \hat{E}_{\text{PN}}(\hat{R}), \end{aligned} \quad (4.24)$$

where

$$\begin{aligned} \hat{E}_{\text{PN}} = & -\frac{5(5-n)}{18} \hat{W}^2 (\tilde{\lambda}_n + n + 7) + \frac{2(5-n)}{3(n+1)} \hat{U}_i \hat{W} (\tilde{\lambda}_n + n + 1) \\ & + \frac{1}{\hat{R}} \left(-\frac{55}{6} \hat{W} + \frac{5}{2} \hat{U}_i \right) - \frac{7}{32\hat{R}^2} \\ & + \frac{\kappa_n \hat{W}}{15\hat{R}^3} \left\{ q_{11} \left(5n - 20 + \frac{5n}{n+1} \gamma_n - \frac{25}{2} \gamma_n \right) + q_s \left(35\gamma_n + 5(5-n) - \frac{20(5-n)}{3} \frac{\eta_n}{\kappa_n} \right) \right\} \\ & + \frac{\kappa_n}{240\hat{R}^4} (302 + 59\alpha_2^2 - 209\alpha_3^2) + O(R^{-5}). \end{aligned} \quad (4.25)$$

For the incompressible case $n = 0$, $\kappa_0 = 1$, $\sigma_0 = 2/35$, $\gamma_0 = 1/7$, $\eta_0 = 6/35$, $\lambda_0 = 8/35$ (see Table I), $U_i = 0$, and $W/M = 2\pi\rho A_0/5$, the expression of E_{PN} , of course, agrees with that obtained in a previous paper [1].

The angular momentum for each star $J_1 = J_N + J_{\text{PN}}/c^2$ is calculated from

$$\begin{aligned} J_N &= \int \rho v_\varphi d^3x, \\ J_{\text{PN}} &= \int \rho \left[v_\varphi \left(\Omega_N^2 \varpi^2 + 6U + \Gamma\varepsilon \right) + \beta_\varphi \Omega \right] d^3x, \end{aligned} \quad (4.26)$$

where $v_\varphi = \Omega \varpi^2$. Using $J_1 \sim J_5$, they are soon expressed as

$$\begin{aligned} J_N &= \Omega \left(\frac{MR^2}{4} + I_s \right) = M^{3/2} a_1^{1/2} \hat{\Omega} \left(\frac{\hat{R}^2}{4} + \frac{\kappa_n q_s}{5} \right) \equiv M^{3/2} a_1^{1/2} \hat{\Omega} \hat{J}_N(\hat{R}), \\ J_{\text{PN}} &= \Omega_N \left[\frac{R^2}{4} (U_i + 5W) + \frac{5}{2} RM^2 + I_s \frac{W}{M} \left\{ \frac{35\gamma_n}{3} + 2(5-n) - \frac{20}{9}(5-n) \frac{\eta_n}{\kappa_n} \right\} \right. \\ &\quad \left. + \frac{M}{4R} (20I_{11} + 15I_{22} - 11I_{33}) \right] \\ &= M^{5/2} a_1^{-1/2} \hat{\Omega}_N \left[\frac{\hat{R}^2}{4} (\hat{U}_i + 5\hat{W}) + \frac{5}{2} \hat{R} + \frac{\kappa_n q_s \hat{W}}{5} \left\{ \frac{35\gamma_n}{3} + 2(5-n) - \frac{20}{9}(5-n) \frac{\eta_n}{\kappa_n} \right\} \right. \\ &\quad \left. + \frac{\kappa_n}{20\hat{R}} (20 + 15\alpha_2^2 - 11\alpha_3^2) \right] \\ &\equiv M^{5/2} a_1^{-1/2} \hat{\Omega}_N \hat{J}_{\text{PN}}(\hat{R}). \end{aligned} \quad (4.27)$$

Once we determine $\alpha_2(\hat{R})$ and $\alpha_3(\hat{R})$ from Eqs.(3.15), we can calculate $\hat{\Omega}_N$, $\hat{\Omega}_{\text{PN}}$, $\hat{E}_N$, $\hat{E}_{\text{PN}}$, $\hat{J}_N$, and $\hat{J}_{\text{PN}}$ as a function of $\hat{R}$ right now using the numerical values for ξ_1 , $|\theta'|$, k_1 , κ_n , σ_n , λ_n and η_n shown in Table I. To evaluate the angular velocity, the energy and the angular momentum, however, we further need the equation for determination of a_1 . We will derive it in the next section.

V. SCALAR VIRIAL RELATION IN THE POST-NEWTONIAN APPROXIMATION

Up to this section, we have derived the equations required for determining the equilibrium sequence except for a_1 (or ρ_c). Here, we derive the equation for a_1 (or ρ_c) in the PN order. Such an equation should come from the scalar virial relation in the PN approximation, which is derived by

$$\begin{aligned}
& \sum_i \int d^3x \rho x_i \frac{\partial}{\partial x_i} \left[K' \rho^{\Gamma-1} - \frac{1}{2c^2} \left(K' \rho^{\Gamma-1} \right)^2 \right] \\
&= \sum_i \int d^3x \rho x_i \frac{\partial}{\partial x_i} \left[U - \frac{X_0}{c^2} + \left\{ \frac{\varpi^2}{2} + \frac{1}{c^2} (2\varpi^2 U - X_\Omega + \beta_\varphi) \right\} \Omega^2 + \frac{\varpi^4}{4c^2} \Omega^4 \right]. \quad (5.1)
\end{aligned}$$

Using the equations shown in previous sections, the left-hand side of Eq.(5.1) can be written as

$$M \left[-\frac{3}{n} k_1 K \rho_c^{1/n} + \frac{1}{c^2} \frac{3(n+1)^2}{n+2} \lambda_n K^2 \rho_c^{2/n} \right]. \quad (5.2)$$

In the right-hand side of Eq.(5.1), many terms are soon integrated as

$$\begin{aligned}
& -W + \frac{3M}{R^3} \mathcal{I}_{11} + \Omega^2 I_s + \frac{1}{c^2} \left[-K_1 + \left(\frac{15+n}{R^3} W + \frac{6M^2}{R^4} \right) \mathcal{I}_{11} \right. \\
& \quad + \Omega_N^2 \left\{ 2 \sum_i (\mathcal{M}_{ii11} + \mathcal{M}_{ii22}) - \frac{R^2}{2} W + 4(I_{U11} + I_{U22}) - K_2 + K_3 \right. \\
& \quad \left. \left. + \frac{M}{8R} (2I_{11} + 15I_{22} - 15I_{33}) \right\} + \frac{\Omega_N^4}{4} R^2 (3I_{11} + I_{22}) + O(R^{-5}) \right], \quad (5.3)
\end{aligned}$$

but for

$$K_1 = \sum_i \int \rho X_{0,i}^{1 \rightarrow 1} x_i d^3x, \quad (5.4)$$

$$K_2 = \sum_i \int \rho X_{\Omega,i}^{1 \rightarrow 1} x_i d^3x, \quad (5.5)$$

$$K_3 = \sum_i \int \rho \beta_{\varphi,i}^{1 \rightarrow 1} x_i d^3x. \quad (5.6)$$

K_1 is evaluated as

$$\begin{aligned}
K_1 &= - \int \int d^3x d^3y \rho(x) \rho_X^{1 \rightarrow 1}(y) \sum_i x_i \frac{\partial}{\partial x_i} \left(\frac{1}{|\mathbf{x} - \mathbf{y}|} \right) \\
&= \int \int d^3x d^3y \rho(x) \rho_X^{1 \rightarrow 1}(y) \sum_i x_i \frac{\partial}{\partial y_i} \left(\frac{1}{|\mathbf{x} - \mathbf{y}|} \right) \\
&= \int d^3y \rho_X^{1 \rightarrow 1}(y) \sum_i D_{i,i} \\
&= \int d^3y \rho_X^{1 \rightarrow 1}(y) (U^{1 \rightarrow 1} + \sum_i y_i U_{,i}^{1 \rightarrow 1}) \\
&= \left(1 + \frac{3}{n} \right) (J_1 + J'_1) + 2(J_2 + J'_2 + J_4 + J'_4), \quad (5.7)
\end{aligned}$$

where we define

$$J'_1 = \sum_i \int \rho U_{,i}^{1 \rightarrow 1} x_i \varepsilon d^3x, \quad (5.8)$$

$$J'_2 = \sum_i \int \rho U_{,i}^{1 \rightarrow 1} x_i U^{1 \rightarrow 1} d^3x, \quad (5.9)$$

$$J'_4 = \sum_i \int \rho U_{,i}^{1 \rightarrow 1} x_i U^{2 \rightarrow 1} d^3x. \quad (5.10)$$

In the ellipsoidal approximation, they are written as

$$\begin{aligned} J'_1 &= -\frac{\tilde{\lambda}_n \xi_1 |\theta'|}{n+2} W K n \rho_c^{1/n}, \\ J'_2 &= -\frac{5-n}{3(n+2)} \frac{W^2}{M} (\tilde{\lambda}_n + n + 2), \\ J'_4 &= -W \left(\frac{M}{R} + \frac{3\mathcal{F}_{11}}{2R^3} + \frac{25\gamma_n}{2(n+1)} \frac{\mathcal{F}_{11}}{R^3} \right) + O(R^{-5}). \end{aligned} \quad (5.11)$$

Using the same method as that adopted in calculating K_1 ,

$$\begin{aligned} K_2 &= 2 \int d^3x \rho \left[\left(x_1^2 + x_2^2 + \frac{R^2}{4} \right) \left(U^{1 \rightarrow 1} + \sum_i x_i U_{,i}^{1 \rightarrow 1} \right) \right] \\ &= \frac{R^2}{2} W + 2(I_{U11} + I_{U22}) + 2 \sum_i (\mathcal{M}_{ii11} + \mathcal{M}_{ii22}). \end{aligned} \quad (5.12)$$

K_3 is rewritten as

$$\begin{aligned} K_3 &= -\frac{1}{8} \int d^3x \rho \sum_i \left[R^2 x_i \frac{\partial}{\partial x_i} \left(7U^{1 \rightarrow 1} + D_{2,2} - U_{,2}^{1 \rightarrow 1} x_2 \right) \right. \\ &\quad \left. + 4x_i \frac{\partial}{\partial x_i} \left(7x_1 D_1 + 7x_2 D_2 + x_1^2 D_{2,2} + x_2^2 D_{1,1} - x_1 x_2 D_{1,2} - x_1 x_2 D_{2,1} \right) \right]. \end{aligned} \quad (5.13)$$

Using the same method as that in calculating J_5 with Eq.(4.19),

$$K_3 = \frac{R^2}{8} (7W - \mathcal{M}_{22}) - \frac{20}{9} \frac{W}{M} I_s \left[\frac{3(5-n)}{5} + \frac{3(n-4)}{n+1} \gamma_n - \frac{\eta_n}{\kappa_n} (5-n) \right]. \quad (5.14)$$

Gathering these results, the PN scalar virial relation is written as

$$\frac{3k_1}{n} K \rho_c^{1/n} + \frac{1}{c^2} B K^2 \rho_c^{2/n} = C \frac{M}{a_1} + D \frac{M^2}{c^2 a_1^2}, \quad (5.15)$$

where

$$\begin{aligned} B &= -\frac{3(n+1)^2}{n+2} \lambda_n, \\ C &= \hat{W} - \frac{\kappa_n q_{11}}{5\hat{R}^3} - \frac{\hat{\Omega}_N^2 \kappa_n q_s}{5} = \frac{3}{n} \hat{U}_i, \end{aligned}$$

$$\begin{aligned}
D = & \frac{(n+3)(5-n)}{3(n+1)n} \hat{W} \hat{U}_i \left(\frac{n-1}{n+2} \tilde{\lambda}_n + n+1 \right) + \frac{2(5-n)}{9} \hat{W}^2 \left(\frac{n-1}{n+2} \tilde{\lambda}_n + n+4 \right) \\
& + \frac{13}{6\hat{R}} \hat{W} + \frac{\kappa_n \hat{W}}{15\hat{R}^3} \left[q_{11} \left(-\frac{11}{2} - 2n + \frac{5(n-4)}{n+1} \gamma_n \right) \right. \\
& \quad \left. - \frac{4}{3} q_s \left(\frac{135-15n}{n+1} \gamma_n + \frac{9n+15}{2} + \frac{10\eta_n}{\kappa_n} (5-n) \right) \right] \\
& - \frac{\kappa_n}{60\hat{R}^4} (86 + 35\alpha_2^2 - 68\alpha_3^2). \tag{5.16}
\end{aligned}$$

Using Eqs.(3.3) and (5.15), $a_1 = a_{1N} + \delta a_1/c^2$ is determined as

$$\begin{aligned}
a_{1N} &= \left(\frac{\hat{U}_i}{k_1 K} \right)^{n/(n-3)} \left(\frac{\xi_1}{4\pi\alpha_2\alpha_3|\theta'|} \right)^{1/(3-n)} M_N^{(n-1)/(n-3)}, \\
\delta a_1 &= a_{1N} \left[\frac{1-n}{3-n} \frac{\delta M}{M_N} + \frac{n}{C(n-3)} \frac{M_N}{a_{1N}} \left\{ D - B \left(\frac{\hat{U}_i}{k_1} \right)^2 \right\} \right], \tag{5.17}
\end{aligned}$$

where we split M as $M_N + \delta M/c^2$. The first term of the equation of δa_1 originates from the Newtonian property, and the second and third terms(which include B and D) are purely the PN origin. Since $B < 0$ and $D > 0$ for $0 < n < 3$, the PN effect tends to decrease δa_1 for realistic neutron stars, and as a result makes the coordinate radius of the star smaller than that in the Newtonian case.

VI. ANALYSIS OF EQUILIBRIUM SEQUENCES

An equilibrium sequence of the compressible Darwin ellipsoids is determined fixing M_* and K . The procedure is as follows.

(1) using Eqs.(3.15), we calculate the equilibrium sequence in the Newtonian order. Up to this stage, $\alpha_2(\hat{R})$ and $\alpha_3(\hat{R})$ are determined, and the results are checked by comparing those by LRS [2].

(2) M is determined by the condition $M_* = \text{constant}$. From Eq.(3.12),

$$M = M_* \left[1 - \frac{M_*}{c^2 a_{1N}} \left\{ 6\hat{W} + \frac{13}{4\hat{R}} + \frac{\kappa_n}{20\hat{R}^3} (15q_{11} + 4q_s) \right\} \right]. \tag{6.1}$$

Then, we set $M_N = M_* = \text{constant}$ and

$$\delta M = -M_* \frac{M_*}{a_{1N}} \left\{ 6\hat{W} + \frac{13}{4\hat{R}} + \frac{\kappa_n}{20\hat{R}^3} (15q_{11} + 4q_s) \right\}. \tag{6.2}$$

(3) after substituting Eqs.(5.17) and (6.1) into the equations for Ω^2 , E_1 and J_1 , we rewrite them as

$$\Omega^2 = \frac{M_*}{a_{1N}^3} \left[\hat{\Omega}_N^2 + \frac{1}{c^2} \left\{ \left(\frac{\delta M}{M_*} - \frac{3\delta a_1}{a_{1N}} \right) \hat{\Omega}_N^2 + \frac{M_*}{a_{1N}} \hat{\Omega}_{PN}^2 \right\} \right], \quad (6.3)$$

$$E_1 = \frac{M_*^2}{a_{1N}} \left[\hat{E}_N + \frac{1}{c^2} \left\{ \left(\frac{2\delta M}{M_*} - \frac{\delta a_1}{a_{1N}} \right) \hat{E}_N + \frac{M_*}{2a_{1N}} \hat{\Omega}_{PN}^2 \left(\frac{\hat{R}^2}{4} + \frac{\kappa_n q_s}{5} \right) + \frac{M_*}{a_{1N}} \hat{E}_{PN} \right\} \right], \quad (6.4)$$

$$J_1 = M_*^{3/2} a_{1N}^{1/2} \hat{\Omega}_N \left[\hat{J}_N + \frac{1}{c^2} \left\{ \left(\frac{3\delta M}{2M_*} + \frac{\delta a_1}{2a_{1N}} + \frac{M_*}{a_{1N}} \frac{\hat{\Omega}_{PN}^2}{2\hat{\Omega}_N^2} \right) \hat{J}_N + \frac{M_*}{a_{1N}} \hat{J}_{PN} \right\} \right], \quad (6.5)$$

and define the total energy and the total angular momentum of the system as $E = 2E_1$ and $J = 2J_1$. Then, using $\alpha_2(\hat{R})$ and $\alpha_3(\hat{R})$ obtained at (1), we calculate the PN sequences of Ω , E , and J as a function of $\hat{R}$.

We repeat these procedures changing the compactness of each star of binary defined as $C_s = M_*/c^2 a_*$, where a_* is the radius of the Newtonian spherical star

$$a_* = M_*^{(n-1)/(n-3)} \xi_1 \left(\frac{n+1}{4\pi} K \right)^{n/(3-n)} (4\pi \xi_1^2 |\theta'|)^{-(1-n)/(3-n)}. \quad (6.6)$$

For the following, we also define

$$\rho_* = \frac{3M_*}{4\pi a_*^3}. \quad (6.7)$$

In figs.1 and 2, we show the energy($E/(M_*^2/a_*)$), the angular momentum($J/\sqrt{M_*^3 a_*}$), and the angular velocity($\Omega/\sqrt{\pi \rho_*}$) as a function of the orbital separation, R_*/a_* for $n = 0.5$ (figs.1) and 1 (figs.2), respectively. In figs.1, the solid and dotted lines denote the results calculated by the ellipsoidal approximation for $C_s = 0$ and 0.02 , and the open and filled circles denote the results by the numerical calculation [4] for $C_s = 0$ and 0.02 , respectively. In figs.2, the solid and dotted lines denote the results calculated by the ellipsoidal approximation for $C_s = 0$ and $1/60$, and the open and filled circles denote the results by the numerical calculation for $C_s = 0$ and $1/60$, respectively. These figures indicate that the ellipsoidal approximation works fairly well for calculating the angular momentum and the angular velocity, but does not very well for calculating the energy: The energy calculated by the ellipsoidal approximation systematically deviates from the numerical results. This is

because we do not rigidly take into account the change of the density profile of the star due to the PN gravity even for $R \rightarrow \infty$, although the energy is sensitive to the density configuration (i.e., $\hat{W}$ or $\hat{A}_0$). The angular momentum and the angular velocity are not sensitive to the density profile, but to the global quantities such as the mass and the spin as it can be found from their equations. However, the deviation of the energy is systematic, so that this weak point could be improved if we artificially reduce $\hat{A}_0$ by a small correction factor which will depend on C_s and $n^\dagger$.

The ellipsoidal approximation is not quantitatively a sophisticated model for calculating the various quantities, in particular the energy, for close binaries of $\hat{R} \sim 3$. Nevertheless, it is a qualitatively good model: Figs.1 and 2 indicate that the coordinate separation at the ISCCO becomes small due to the PN effect[‡], and this feature agrees with the previous numerical results [4]. We show the relation between the angular momentum and the angular velocity for $n = 0.5$ and 1.0 in figs.3. These indicate that Ω at the ISCCO increases with the PN effect, and this feature also agrees with the numerical results [4].

In figs.4–8, we show the angular momentum as a function of R_* (a) and the angular velocity as a function of the angular momentum(b) for $n = 0.25$ (figs.4), 0.5 (figs.5), 0.75 (figs.6), 1.0 (figs.7), 1.5 (figs.8) and for $C_s = 0$ (solid line), 0.01 (dotted line), 0.02 (dashed line), 0.03 (long dashed line), respectively. We can see the feature found in figs.1–3 again; i.e., (1) the coordinate radius at the ISCCO decreases with the increase of C_s , and (2) the orbital angular velocity at the ISCCO increases with C_s . Thus, we conclude that such fea-

[†] For the same C_s , we will need a larger correction factor for larger n , and for $n = 0$, we do not need the correction.

[‡] Since we use the ellipsoidal approximation for the density profile which is not the strict one, and also neglect sufficiently high order terms of R^{-1} in the energy and the angular momentum, the location of the energy minimum does not coincide with that of the angular momentum minimum. However, the difference is not large like the results in a previous paper [1].

tures hold irrespective of the EOS if the ISCCO exists at the Newtonian order. These figures also indicate that the star of a softer EOS is more susceptible to the PN effect, and as a result the orbital radius and the angular velocity at the ISCCO change much.

Finally, we argue the stability of each star against the collapse due to the influence of the GR gravity from the companion star considering the central density ρ_c . From Eqs.(3.3), (5.17), and (6.1), we obtain

$$\begin{aligned}\rho_c &= \rho_N \left[1 + \frac{1}{c^2} \left(\frac{\delta M}{M_*} - \frac{3\delta a_1}{a_{1N}} \right) \right] \\ &= \rho_N \left[1 + \frac{M_*}{c^2 a_{1N}} \left\{ \frac{n}{3(3-n)} \left(19 + 4n - 3n^2 + \frac{(3n-1)(5-n)\tilde{\lambda}_n}{n+1} \right) \hat{W} \right. \right. \\ &\quad \left. \left. + \frac{A_3}{\hat{R}^3} + \frac{A_4}{\hat{R}^4} + O(\hat{R}^{-5}) \right\} \right],\end{aligned}\tag{6.8}$$

where

$$\rho_N = \frac{M_* \xi_1}{4\pi |\theta'| \alpha_2 \alpha_3 a_{1N}^3},\tag{6.9}$$

and A_3 and A_4 are functions of $\hat{R}$. Note that in $\hat{W}$, the effect of $O(\hat{R}^{-3})$ is implicitly included because α_2 and α_3 behave as $1 + O(\hat{R}^{-3})$. The lowest order terms of $\hat{R}$ (the terms of $O(\hat{R}^0)$) are concerned in the well known GR instability to the radial collapse for each star itself [10]: Since the coefficient of $\hat{W}$ is always positive for $0 < n < 3$, in the case of $n \rightarrow 3$ or of sufficiently large $\hat{W}$, $\rho_c a_*^2 / c^2 \sim M_* / (c^2 a_*)$ becomes larger than unity, which means that the star is swallowed inside its Schwarzschild radius to be a black hole due to the GR gravity.

In Eq.(6.8), the terms of $O(R^{-1})$ do not appear: As Wiseman points out, this is the consequence of the strong equivalence principle [11]. Thus, the effect due to the GR gravity of the companion star appears from the terms of $O(R^{-3})$ which are concerned in the tidal effect in the PN order. As Lai pointed out [12], the Newtonian tidal effect tends to stabilize each star; i.e., it reduces the central density. Actually, $\partial \rho_N / \partial R$ is always positive in the present case. Here, we demonstrate that the PN tidal effect also reduces the central density. In figs.9, we show ρ_c / ρ_N as a function of the orbital separation for $n = 0.5$ (a), 1(b), and 1.5(c). In each figure, the solid, dotted, dashed, and long dashed lines denote the cases for

$C_s = 0, 0.01, 0.02$ and 0.03 , respectively. These figures indicate that the GR effect due to the companion star reduces the central density. This seems to mean that it will act to stabilize each star of binary. Recently, Wilson et al. calculated the equilibrium state of binary neutron stars using the semi-relativistic approximation [13], and they showed that the central density of each star increases with the decrease of the orbital separation for a critical orbital separation. However, the present analysis suggests that for the corotating binary neutron stars, such an effect cannot be explained by the first PN correction of general relativity.

VII. SUMMARY

Using the ellipsoidal model for the density configuration (the ellipsoidal approximation), we have calculated the equilibrium states of corotating binary stars of the stiff polytropic EOS in the first PN approximation of general relativity. Comparing the equilibrium sequences with those obtained by numerical calculation [4], we confirm that the ellipsoidal approximation is a qualitatively good model to see the several PN effects in binary neutron stars, and then get the conclusions as follows.

- (1) Irrespective of the polytropic index n , the PN effects make the orbital separation at the ISCCO smaller and the orbital angular velocity at the ISCCO larger.
- (2) As in the Newtonian case [12], the PN tidal effect from the companion star reduces the central density of each star. Thus, it seems to act the stabilization of each star.

These results will be helpful to check numerical results in PN and/or full GR numerical computations of obtaining the equilibrium states of binary neutron stars.

ACKNOWLEDGMENTS

Numerical computations were partly performed on FACOM VPP-300 in data processing center of National Astronomical Observatory in Japan. This work was in part sup-

ported by Japanese Grant-in-Aid of Ministry of Education, Culture, Science and Sports
(Nos. 08237210 and 08NP0801).

REFERENCES

- [1] K. Taniguchi and M. Shibata, submitted to Phys. Rev. D.
- [2] D. Lai, F. A. Rasio, and S. L. Shapiro, *Astrophys. J. Supple.* **88**, 205(1993); *Astrophys. J.* **420**, 811(1994).
- [3] For example, S. L. Shapiro and S. A. Teukolsky, *Black Holes, White Dwarfs, and Neutron Stars*(New York, Wiley, 1983).
- [4] M. Shibata, submitted to Phys. Rev. D;
M. Shibata, *Prog. Theor. Phys.* **96**, 317(1996).
- [5] S. Chandrasekhar, *Astrophys. J.* **142**, 1488(1965).
- [6] S. Chandrasekhar, *Astrophys. J.* **148**, 621(1967), and also see nice collection, *Selected Papers*, S. Chandrasekhar, Vol.5 (Chicago University Press, 1990).
- [7] H. Asada and M. Shibata, *Phys. Rev. D* **54**, 4944(1996).
- [8] S. Chandrasekhar, *Ellipsoidal Figures of Equilibrium* (Yale University Press, New Haven, 1969).
- [9] C. M. Will, *Theory and Experiment in Gravitational Physics* (Cambridge University Press, Cambridge, 1981), p.146.
- [10] S. Chandrasekhar, *Astrophys. J.* **140**, 417(1964); **142**, 1519(1965).
- [11] A. G. Wiseman, gr-qc 9704018.
- [12] D. Lai, *Phys. Rev. Lett.* **76**, 4878(1996): He showed that the Newtonian tidal effect acts to stabilize each star of binary.
- [13] J. R. Wilson, G. J. Mathews, and P. Marrinetti, *Phys. Rev. D.* **54**, 1317(1996).

Table I

Quantities for the polytropic stars of different polytropic indices.

n	ξ_1	$ \theta' $	k_1	κ_n	σ_n	λ_n	η_n
0	$\sqrt{6}$	$2/\sqrt{6}$	0	1	2/35	8/35	6/35
0.25	2.59209	.633717	.108069	.903969	.0561363	.252922	.156543
0.5	2.75270	.499997	.229390	.814828	.0538683	.273988	.143283
0.75	2.93452	.398094	.360772	.731532	.0507645	.292471	.131209
1.0	3.14159	.318310	.500000	.653455	.0471121	.308844	.120046
1.5	3.65375	.203301	.795872	.511500	.0389291	.336503	.0997846

Figure captions

Figs.1 The energy($E/(M_*^2/a_*)$, (a)), the angular momentum($J/\sqrt{M_*^3 a_*}$, (b)), and the angular velocity($\Omega/\sqrt{\pi\rho_*}$, (c)) as a function of the orbital separation(R_*/a_*) for $n = 0.5$. The solid and dotted lines denote the results calculated by the ellipsoidal approximation for $C_s = 0$ and 0.02, and the open and filled circles denote the results by the numerical calculation for $C_s = 0$ and 0.02, respectively. Note that the innermost point of the sequence is determined from the condition $2a_1 = R$.

Figs.2 The same as in figs.1, but for $n = 1$ and $C_s = 0, 1/60$.

Figs.3 The relation between the angular momentum and the angular velocity for $n = 0.5$ (a) and 1.0(b). The meaning of the lines and marks is the same as that in figs.1 or 2.

Figs.4 The angular momentum as a function of R_*/a_* (a) and the angular velocity as a function of the angular momentum (b) for $n = 0.25$, and for $C_s = 0$ (solid line), 0.01(dotted line), 0.02(dashed line), 0.03(long dashed line).

Figs.5 The same as figs.5, but for $n = 0.5$.


Figs.6 The same as figs.5, but for $n = 0.75$.

Figs.7 The same as figs.5, but for $n = 1.0$.


Figs.8 The same as figs.5, but for $n = 1.5$.

Figs.9 ρ_c/ρ_N as a function of the orbital separation for $n = 0.5$ (a), 1.0(b), and 1.5(c). In each figure, the solid, dotted, dashed, and long dashed lines denote the results for $C_s = 0, 0.01, 0.02$ and 0.03, respectively.


$n=0.5$



(a)






(b)



(c)

Figs.1



Figs.2









Fig.3





Figs.4

n=0.5




Figs.5

$n=0.75$





Figs.6




Figs.7

$n=1.5$



Figs.8



Figs.9